
PALOS VERDES PENINSULA SUMMER SCHOOL Health Services

SEIZURE EMERGENCY PLAN


Student's Name:		DOB:	Grade:		
Current Medication(s):					
Type of Seizure(s)	<u>Description</u> *				
☐ Absence	Staring, eye blinking, l				
☐ Simple partial	Remains conscious, distorted sense of smell, hearing, sight, involuntary rhythmic jerking/twitching on one side, Other				
☐ Complex partial	Confused, not fully responsive/unresponsive, may appear fearful, purposeless, repetitive movements, Other				
☐ Generalized tonic-clonic	•	g, breathing may be shallow, lip ousness, confusion, weariness			
*Student may experience some or all of	the listed symptoms during a	a seizure.			
Possible warning and/or behavior changes prior to the seizure					
Average Frequency:					
Usual Time of Day Seizure Occurs:					
Average Length of time Seizure Lasts:					
Two rago congin of time concare cases.					
TREATMENT □ DIASTAT (diazepam rectal gel)mg rectally prn for: seizure >minutes OR foror more seizures inhours □ Use VNS (vagal nerve stimulator) Magnet					
Other					
□ CALL 911 if: -Seizure does not stop by its -Seizure does not stop withir -Child does not start waking -Child does not start waking	ıminutes of gi up withinminu		IASTAT given) DIASTAT is given)		

Foll	 Owing a Seizure: Child should rest in healt Child may return to class Parents should be notified 			
Phy		Tiote Sent nome with the child	Date	
•	ense#			
Parent's Signature:				
EM	IERGENCY CONTACTS	3		
1.	Name:			
	Phone:	Relationship:		
2.	Name:			
	Phone:	Relationship:		
3.	Name:			
	Phone:	Relationship:		
TR	AINED STAFF MEMBE	RS		
1.	Name:		Room:	
2.	Name:		Room:	
3.	Name:		Room:	

First Aid for Seizures









In case of seizure, immediately contact the school nurse or designed

Cushion head, remove glasses

Loosen tight clothing








Turn on side and keep airway clear

Note the time a seizure starts and the length of time it lasts

Don't put anything in mouth







Don't hold down

As seizure ends...

...offer help

Most seizures in people with epilepsy end without harm after 1 or 2 minutes and usually do not require a trip to the emergency room. But sometimes there are good reasons to call for emergency help:

- Prolonged seizure
- Slow recovery
- Repeated seizures on the same day
- Difficulty breathing after a seizure
- Pregnancy or other medical diagnosis
- Any signs of injury

CONFIDENTIALITY STATEMENT

By creating and distributing the DIASTAT® (diazepam rectal gel) Back-to-School Kit, it is the expressed intention of Xcel Pharmaceuticals, Inc. to provide a value-added service to healthcare providers, caregivers, and patients. Xcel Pharmaceuticals, Inc. intends to fully comply with HIPAA guidelines for patient confidentiality to ensure that personal information is not disclosed to anyone outside of those that the parent/caregiver, healthcare professional, or student-patient deem acceptable. Xcel Pharmaceuticals, Inc. is not responsible for the DIASTAT Back-to-School Kit or its contents after the kit has been released to healthcare professionals. Careful consideration should be given before any medical information is released to others.

DIASTAT is a registered trademark of Xcel Pharmaceuticals, inc.

® 2004 Xcel Pharmaceuticals, inc.

